ALKA-SELTZER PLUS MAXIMUM STRENGTH SEVERE SINUS CONGESTION AND COUGH DAY AND NIGHT- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride Bayer HealthCare LLC.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Day & Night Liquid Gels

Do not take these products at the same time.

Drug Facts

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Day Liquid Gels

Active ingredients (in each capsule)

Acetaminophen 325 mg

Dextromethorphan hydrobromide 10 mg

Phenylephrine hydrochloride 5 mg

Purposes

Pain reliever/fever reducer Cough suppressant

Nasal decongestant

Uses

- · temporarily relieves these symptoms due to a cold or flu:
- · nasal congestion · sinus congestion and pressure
- · headache · minor aches and pains
- · cough · sore throat
- · helps clear nasal passages and shrinks swollen membranes
- · temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- · more than 4,000 mg of acetaminophen in 24 hours
- · with other drugs containing acetaminophen
- · 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe

allergic reactions. Symptoms may include:

- · skin reddening · blisters · rash · hives
- · facial swelling · asthma (wheezing) · shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than

2 days, is accompanied or followed by fever, headache, rash, nausea,

or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease heart disease high blood pressure
- thyroid disease diabetes
- cough with excessive phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin **When using this product** do not exceed recommended dosage **Stop use and ask a doctor if**

- \cdot pain, cough, or nasal congestion gets worse or lasts more than 7 days
- · fever gets worse or lasts more than 3 days
- · redness or swelling is present
- · new symptoms occur
- · cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

· nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Directions

· do not take more than the recommended dose

- · adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- · children under 12 years: do not use

Other information

Other information

· store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients FD&C yellow #6, FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments?

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM –5PM EST)

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Night Liquid Gels Drug Facts

Active ingredients (in each capsule)

Acetaminophen 325 mg

Dextromethorphan hydrobromide 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine hydrochloride 5 mg

Uses

- · temporarily relieves these symptoms due to a cold or flu:
- · nasal congestion · sinus congestion and pressure
- · headache · minor aches and pains
- · cough · sore throat
- · runny nose · sneezing
- · helps clear nasal passages and shrinks swollen membranes
- · temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- · more than 4,000 mg of acetaminophen in 24 hours
- · with other drugs containing acetaminophen
- · 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe

allergic reactions. Symptoms may include:

· skin reddening · blisters · rash · hives

· facial swelling · asthma (wheezing) · shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than

2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Ask a doctor before use if you have

- liver disease heart disease high blood pressure
- thyroid disease diabetes glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- · pain, cough, or nasal congestion gets worse or lasts more than 7 days
- · fever gets worse or lasts more than 3 days
- · redness or swelling is present
- · new symptoms occur
- · cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

· nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help

or contact a Poison Control Center right away. Quick medical attention

is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- · do not take more than the recommended dose
- · adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- · children under 12 years: do not use

Other information

Other information


 \cdot store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue #1, D&C yellow #10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments?

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)





37/8

Cut Edge

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Day Liquid Gels Sinus Congestion & Cough Night Liquid Gels

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MACI) (certain drugs for degression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MACI drug. If you do not know if your prescription drug contains an MACI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
 heart disease
 high blood pressure
- th vroid disease
 diabetes

asthma, or emphysema

- cough with excessive phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
 persistent or chronic cough such as occurs with smoking,

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nongrescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MACI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MACI drug. If you do not know if your prescription drug combins an MACI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age


Ask a doctor before use if you have

- li ver disease
 heart disease
 high blood pressure
- thyroid disease
 diabetes
 - oetes glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
 provide text or charging equals such as procure with excepting.
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

INSIDE of EC Flap



ALKA-SELTZER PLUS MAXIMUM STRENGTH SEVERE SINUS CONGESTION AND COUGH DAY AND NIGHT

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride kit

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0280-1610

Packaging

Item Code Package Description Marketing Start Date Marketing End Date
1 NDC:0280-1610-20 1 in 1 CARTON; Type 0: Not a Combination Product 06/12/2017

Quantity of Parts

Quality of Faits			
Part # Package Quantity Total Product	t Quantity		
Part 1 2 BLISTER PACK 12			
Part 2 4 BLISTER PACK 16			

Part 1 of 2

ALKA-SELTZER PLUS MAXIMUM STRENGTH SEVERE SINUS CONGESTION AND COUGH DAY

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	325 mg		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg		
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg		

Inactive Ingredients				
Ingredient Name	Strength			
SORBITOL (UNII: 506T60A25R)				
WATER (UNII: 059QF0KO0R)				
SHELLAC (UNII: 46N107B71O)				
SORBITAN (UNII: 6092ICV9RU)				
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GLYCERIN (UNII: PDC6A3C0OX)				
GELATIN (UNII: 2G86QN327L)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
PO VIDO NE (UNII: FZ989 GH94E)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SODIUM HYDROXIDE (UNII: 55X04QC32I)				

Product Characteristics			
Color	orange	Score	no score
Shape	OVAL	Size	20 mm
Flavor		Imprint Code	AS;SC
Contains			

Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1		2 in 1 CARTON			
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product			

	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ı	OTC monograph final	nart341	06/12/2017	

Part 2 of 2

ALKA-SELTZER PLUS MAXIMUM STRENGTH SEVERE SINUS CONGESTION AND COUGH NIGHT

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg		
ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	325 mg		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg		
DO XYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DO XYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg		

Inactive Ingredients				
Ingredient Name	Strength			
WATER (UNII: 059QF0KO0R)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
GELATIN (UNII: 2G86QN327L)				
GLYCERIN (UNII: PDC6A3C0OX)				
SHELLAC (UNII: 46 N107B710)				
SO DIUM HYDRO XIDE (UNII: 55X04QC32I)				
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)				
PO VIDO NE (UNII: FZ989 GH94E)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
SORBITOL (UNII: 506T60A25R)				
SORBITAN (UNII: 6092ICV9RU)				

Product Characteristics			
Calan	groon	Canada	no scoro

Shape	OVAL	Size	20 mm
Flavor		Imprint Code	AS;SC
Contains			

Packaging					
# Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	2 in 1 CARTON				
1	4 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	06/12/2017		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	06/12/2017		

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 11/2019 Bayer HealthCare LLC.